This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

three three three real effects have three three

۲. إ

Claim 17, line 1, change "any one of the preceding claims" to --claim 1--.

Claim 19, line 1, change "any one of the preceding claims" to --claim 1--.

Claim 21, lines 1-2, change "any one of the preceding claims" to --claim 1--.

Please cancel Claim 27 without prejudice or disclaimer.

Claim 28, lines 1-2, change "any one of claims 1 to 20" to --claim 1--.

Claim 29, lines 1-2, change "any one of claims 1 to 20" to --claim 1--.

REMARKS

The above amendments are made to reduce initial filing fees by removing multiple dependent claims.

Respectfully submitted,

NIXON & VANDERHYE P.C.

By:

Arthur R. Crawford Reg. No. 25,327

ARC:ms

1100 North Glebe Road, 8th Floor

Arlington, VA 22201-4714 Telephone: (703) 816-4000 Facsimile: (703) 816-4100

Claims

A protein having luciferase activity and at least 60% similarity to luciferase from Photinus pyralis, Luciola mingrelica, Luciola cruciata, Luciola lateralis, Hotaria paroula, Pyrophorus plagiophthalamus, Lampyris noctiluca, Pyrocoelia nayako or Photinus pennsylanvanica, wherein in the sequence of the enzyme, at least one of (a) the amino acid residue corresponding to residue 214 in 10 Photinus pyralis luciferase or to residue 216 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase; (b) the amino acid residue corresponding to residue 232 in Photinus pyralis luciferase or to residue 234 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase; (c) amino acid residue corresponding to residue 295 in Photinus pyralis luciferase or to residue 297 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase; (d) amino acid residue corresponding to amino acid 14 of the Photinus pyralis luciferase or to residue 16 of Luciola mingrelica, or 17 of Luciola cruciata or Luciola lateralis; (e) amino acid residue corresponding to amino acid 35 of the Photinus pyralis luciferase or to residue 37 of Luciola mingrelica, or 38 of Luciola cruciata or Luciola lateralis; (f) the amino acid residue corresponding to amino acid residue 105 of the Photinus pyralis luciferase or to residue 106 of 25 Luciola mingrelica, 107 of Luciola cruciata or Luciola

(g) amino acid residue corresponding to amino acid residue 234 of the *Photinus pyralis* luciferase or to residue 236 of *Luciola mingrelica*, *Luciola cruciata* or *Luciola lateralis*;

lateralis or 108 of Luciola lateralis gene;

- (h) amino acid residue corresponding to amino acid residue 420 of the Photinus pyralis luciferase or to residue 422 of Luciola mingrelica, Luciola cruciata or Luciola lateralis;
- (i) amino acid residue corresponding to amino acid residue 310
 of the Photinus pyralis luciferase or to residue 312 of Luciola mingrelica, Luciola cruciata or Luciola lateralis;

is different to the amino acid which appears in the corresponding wild type sequence and wherein the luciferase enzyme has increased thermostability as compared to an enzyme having the amino acid of the corresponding wild-type luciferase at this position.

2. A protein according to claim 1 which has the sequence of a wild-type luciferase, in which more than one amino acid residue is different to that of the wild type enzyme.

10

5

3. A protein according to claim 2 wherein up to 50 amino acids are different to that of the wild type enzyme.

i

£:ī

ΓIJ

4. A protein according to any one of the preceding slaims wherein the luciferase is a modified form of luciferase of Photinus pyralis, Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase.

- 5. A protein according to any one of the preceding claims wherein the sequence of luciferase of *Photinus pyralis*, wherein at least one of
- (a) the amino acid residue corresponding to residue 214 in *Photinus pyralis* luciferase is other than threonine;
- (b) the amino acid residue corresponding to residue 232 in
- 25 Photinus pyralis luciferase is other than isoleucine;
 - (c) amino acid residue corresponding to residue 295 in *Photinus* pyralis luciferase is other than phenylalanine;
 - (d) amino acid residue corresponding to amino acid 14 of the *Photinus pyralis* luciferase is other than phenylalanine;
- 30 (e) amino acid residue corresponding to amino acid 35 of the Photinus pyralis luciferase is other than leucine;
 - (f) amino acid residue corresponding to amino acid residue 105 of the *Photinus pyralis* luciferase is other than alanine;
 - (g) amino acid residue corresponding to amino acid residue 234
- of the *Photinus pyralis* luciferase is other than aspartic acid; (h) amino acid residue corresponding to amino acid residue 420
 - of the Photinus pyralis luciferase is other than serine;

(i) amino acid residue corresponding to amino acid residue 310 of the *Photinus pyralis* luciferase is other than histidine.

 \mathcal{N}

10

ij

ξÑ

Į.;

îIJ

:: =:=

[]

fij

f ij

[]

ļ.:£

- 6. A protein according to any one of claims 1 to 4 wherein protein has substantially the sequence of Luciola mingrelica, Luciola cruciata or Luciola lateralis enzyme, and wherein at least one of
- (a) the amino acid residue corresponding to residue 216 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase is other than glycine (for Luciola mingrelica based sequences) or aparagine (for Luciola cruciata or Luciola lateralis) based sequences;
- (b) the amino acid residue corresponding to residue 234 of Luciola mingrelica, Luciola cruciata or Luciola lateralis
- 15 luciferase is other than serine;
 - (c) amino acid residue corresponding to residue 297 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase is other than leucine;
 - (d) amino acid residue corresponding to amino acid 16 of Luciola mingrelica, Luciola cruciata or Luciola lateralis is other than phenylalanine;
 - (e) amino acid residue corresponding to residue 37 of Luciola mingrelica, or residue 38 of Luciola cruciata and Luciola lateralis is other than lysine;
 - 25 (f) amino acid residue corresponding to amino acid residue 106 of Luciola mingrelica, 107 of Luciola cruciata or Luciola lateralis, or 108 of Luciola lateralis gene is other than glycine;
 - (g) amino acid residue corresponding to amino acid residue 236
 30 of Luciola mingrelica, Luciola cruciata or Luciola lateralis is other than glycine;
 - (h) amino acid residue corresponding to residue 422 of Luciola mingrelica, Luciola cruciata or Luciola lateralis is other than threonine;
 - 35 (i) amino acid residue corresponding to amino acid residue 312 of Luciola mingrelica, Luciola cruciata or Luciola lateralis is other than threonine (for Luciola mingrelica based sequences)

WO 00/24878 PCT/GB99/03538

or valine (for Luciola cruciata or Luciola lateralis) based sequences.

- 7. A protein according to any one of the preceding claims

 5 wherein comprising a protein having luciferase activity and at least 60% similarity to luciferase from Photinus pyralis,

 Luciola mingrelica, Luciola cruciata or Luciola lateralis
 - (a) the amino acid residue corresponding to residue 214 in

enzyme wherein in the sequence of the enzyme, at least one of

10 Photinus pyralis luciferase and to residue 216 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase is mutated and is other than threonine in the case of Photinus pyralis luciferase; or

£:1

7.1. 7.1.

[1]

fIJ

:: ::=

!!!

711

[]

į.;

30

35

Œ,

- (b) the amino acid residue corresponding to residue 232 in Photinus pyralis luciferase and to residue 234 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase is mutated and is other than isoleucine in the case of Photinus pyralis luciferase; or
- (c) amino acid residue corresponding to residue 295 in *Photinus* pyralis luciferase and to residue 297 of *Luciola mingrelica*, *Luciola cruciata* or *Luciola lateralis* luciferase is mutated and is for example, other than phenylalanine in the case of *Photinus pyralis* luciferase;

and the luciferase enzyme has increased thermostability as compared to the wild-type luciferase.

- 8. A protein according to claim 1 wherein the amino acid residue corresponding to residue 214 in *Photinus pyralis* luciferase and to residue 216 of *Luciola mingrelica*, *Luciola cruciata* or *Luciola lateralis* luciferase is alanine.
- 9. A protein according to any one of the preceding claims wherein the amino acid residue corresponding to residue 232 in Photinus pyralis luciferase and to residue 234 of Luciola mingrelica, Luciola cruciata or Luciola lateralis luciferase is alanine.

WO 00/24878 PCT/GB99/03538

28

10. A protein according to any one of the preceding claims which is a mutated *Photinus pyralis* luciferase wherein the amino acid residue corresponding to residue 295 in *Photinus pyralis* luciferase is leucine.

0

11. A protein according to any one of the preceding claims wherein the amino acid residue corresponding to amino acid 14 of the *Photinus pyralis* luciferase or to amino acid 16 in Luciola luciferase, is alanine.

10

12. A protein according to any one of the preceding claims wherein the luciferase is a mutated luciferase of *Photinus* pyralis or a Luciola species where the amino acid residue corresponding to amino acid 35 of the *Photinus* pyralis luciferase or to amino acid residue 37 in Luciola mingrelica or 38 of Luciola lateralis or cruciata luciferase is alanine.

15 III

13. A protein according to any one of the preceding claims wherein the amino acid residue corresponding to residue 105 in Photinus pyralis luciferase and to residue 106 of Luciola mingrelica, 107 of Luciola cruciata or Luciola lateralis or 108 of Luciola lateralis gene luciferase is valine.

 \mathcal{K}

14. A protein according to any one of the preceding claims which comprises a mutated *Photinus pyralis* luciferase wherein the amino acid residue corresponding to residue 234 in *Photinus pyralis* luciferase is glycine.

€ 30

15. A protein according to any one of the preceding claims which comprises a mutated *Photinus pyralis* luciferase wherein the amino acid residue corresponding to residue 420 in *Photinus pyralis* luciferase is threonine.

€ :

16. A protein according to any one of the preceding claims which comprises a mutated *Photinus pyralis* luciferase wherein the amino acid residue corresponding to residue 310 in *Photinus pyralis* luciferase is arginine.

WO 00/24878 PCT/GB99/03538

29

17. A protein according to any one of the preceding claims wherein the amino acid at position corresponding to amino acid 354 of the *Photinus pyralis* luciferase (356 in Luciola luciferase) is other than glutamate.

5

18. A protein according to claim 17 wherein the amino acid at position corresponding to amino acid 354 of the *Photinus* pyralis luciferase (356 in Luciola luciferase) is lysine or arginine.

10

19. A protein according to any one of the preceding claims wherein the amino acid at the position corresponding to amino acid 217 in Luciola luciferase (215 in *Photinus pyralis*) is a different hydrophobic amino acid.

15

the of the tend

Į.

f

::=

[]

20. A protein according to claim 19 wherein the amino acid at the position corresponding to amino acid 217 in Luciola luciferase (215 in *Photinus pyralis*) is isoleucine, leucine or valine.

20

21. A nucleic acid which encodes a luciferase according to any one of the preceding claims.

25

- 22. A vector comprising a nucleic acid according to claim 21.
- 23. A cell transformed with a vector according to claim 22.
- 24. A cell according to claim 23 which is a prokaryotic cell.
- 30 25. A cell according to claim 23 which is a plant cell.
 - 26. A plant comprising cells according to claim 25.
- 27. A method of producing a protein according to any one of claims 1 to 20, which method comprises culture of a cell according to claim 23 or growth of a plant according to claim 26.

- 28. The use of a protein according to any one of claims 1 to 20 in a bioluminescent assay.
 - 29. A kit comprising a protein according to any one of claims
 - 30. A kit according to claim 29 which further comprises luciferin.

the graph of graph and graph in the second state of the second sta

AFLII-SENSE / 6373

AFLII-ANTI / 6374

Fig.5.

CGCCGCTGAGCTCCCCGCCGCCG

SACI-SENSE / 6371

CGGCGGCGGGAGCTCACCGGCG

SACI-ANTI / 6372

CCTTGTATTTAATTAAAGACTTAAGGCGGTCAACTATGAAGAAGTGTTCG

P14A-SENSE / 6375

F14A-ANTI / 6376 C-TERMLSEQ/ 6641

CCTCTAGCGGATAGGCTGGTGCCGGGCCTTTC CCATAAATTTACCGAATTCGTCGACTTCGATCGAGG **GTGTGGAATTGTGAGCGG**

N-TERMLSEQ / 6651

GAGATACGCCGCGGTTCCTGG L35A-SENSE / 6682 CCAGGAACCGCGGCGTATCTC L35A-SENSE / 6683

CCCTATTTTCATTCCTGGCCAAAAGCACTG GAGTGCTTTTGGCCAGGAATGAAAATAGGG CCGCATAGACTCTCTGCGTCAGATTC GAATCTGACGCAGAGAGCTCTATGCGG

GAAAGGCCCGGCACCAGCCTATCCTCTAGAGG

F295L-SENSE/ 9048 F295L-ANTI / 9049 T214A + A215L-SENSE / 9063 T214A + A215L-ANTI / 9064 Insertion of BamHI at G339 / 9077

GTATAGATTTGAAAAAGAGCTG CAGCTCTTTTTCAAATCTATAC **GGCTACATACTGGAGACATAGC GCTATGTCTCCAGTATGTAGCC**

GTTGACCGCTTGGGATCCTTAATTAAATAC

E270K-SENSE / 257 E270K-ANTI / 258 **\$420T-SENSE / 629** S420T-ANTI / 630

GCAGTTGCGCCCGTGAACGAC A105L-SENSE / 790 GTCGTTCACGGGCGCAACTGC A105L-ANTI /791

CAAATCATTCCGGGTACTGCGATTTTAAG CTTAAAATCGCAGTACCCGGAATGATTTG

D234G-SENSE / 792 D234G-ANTI / 793

CCGCATAGAACTCTCTGCGTCAGATTC A215L-SENSE / 7726 GAATCTGACGCAGAGAGTTCTATGCGC A215L-ANTI / 7727 CTGATTACACCCAAGGGGGATG E354K-SENSE / 7792 CATCCCCTTGGGTGTAATCAG E354K-ANTI / 7793 cccttccgcatagannngcctgcgtcagt

actgacgcaggcNNNtctatgcggaaggg

T214N-Sense / 8202 T214N-Anti / 82033

GCAATCAAATCGCTCCGGATACTGC GCAGTATCCGGAGCGATTTGATTGC

1232A-SENSE / 6911 1232A-ANTI / 6912

CCATTCCATCAAGGTTTTGG

H245Q-SENSE / 9128

CCAAAACCTTGATGGAATGG

H245Q-ANTI / 9129